Physics 2A: Lecture 7

Today's Agenda

- Uniform Circular Motion
- Centripetal acceleration
- Examples
 - Car on track
 - Carnival ride
 - Loop-de-loop



Start Recording!

Clicker Question 1:

Suppose you are driving through a valley whose bottom has a circular shape. If your mass is m, what is the magnitude of the normal force F_N exerted on you by the car seat as you drive past the bottom of the hill?





Clicker 2:

A car that's out of gas coasts over the top of a hill at a steady 20 m/s. Assume air resistance is negligible. Which free-body diagram describes the car at this instant?

A.



 \vec{v}

© 2017 Pearson Education, Inc.

Clicker Question 3:

Harvey is riding a carnival ride which consists of a rotating cylindrically shaped room of radius R = 8 m. Harvey's mass is 85 kg. The floor drops out from below Harvey's feet, and he sticks to the wall rather than descending with the floor. The coefficient of static friction of the wall is $\mu_s = 0.766$.

How many forces act on Harvey?

(a) 1
(b) 2
(c) 3
(d) 4
(e) 5



How many forces act on Harvey?



Clicker Question 4:

Harvey is riding a carnival ride which consists of a rotating cylindrically shaped room of radius R = 8 m. Harvey's mass is 85 kg. The floor drops out from below Harvey's feet, and he sticks to the wall rather than descending with the floor. The coefficient of static friction of the wall is $\mu_s = 0.766$.

What force provides the centripetal acceleration required to keep Harvey moving in a circle?

- (a) Gravity
- (b) Static Friction
- (c) Normal Force



What force provides the centripetal acceleration required to keep Harvey moving in a circle?



What force provides the centripetal acceleration required to keep Harvey moving in a circle?



Student: why is there no force opposing the centripetal acceleration? Shouldnt the forces be balanced or else the person will fly towards the center?

Clicker Question 5:

Harvey is riding a carnival ride which consists of a rotating cylindrically shaped room of radius R = 8 m. Harvey's mass is 85 kg. The floor drops out from below Harvey's feet, and he sticks to the wall rather than descending with the floor. The coefficient of static friction of the wall is $\mu_s = 0.766$.

What are the equations of motion for Harvey?

(a)
$$F_S = \mu_s mg$$
, $F_N = mg$
(b) $F_S = mgsin\theta$, $F_N = mgcos\theta$
(c) $F_S = mg$, $F_N = ma_C$
(d) $F_S = ma_C$, $F_N = mg$
(e) $F_S = mg$, $F_N = 0$



What are the equations of motion for Harvey?



Clicker Question 6:

Harvey is riding a carnival ride which consists of a rotating cylindrically shaped room of radius R = 8 m. Harvey's mass is 85 kg. The floor drops out from below Harvey's feet, and he sticks to the wall rather than descending with the floor. The coefficient of static friction of the wall is $\mu_s = 0.766$.

Harvey's speed must be at least

(a) 0.80 m/s
(b) 2.00 m/s
(c) 3.20 m/s
(d) 5.60 m/s
(e) 10.12 m/s



Harvey's speed must be at least



Clicker Question 7:

A box rests on an incline. The force of gravity pulls straight down as shown. When dealing with inclined planes we will often find it convenient to rotate our coordinate system as shown. How would you express the gravitational force in the new *x*-direction?

- (a) $F_g \cos(\phi)$ (b) $F_g \sin(\phi)$ (c) $-F_g \cos(\phi)$ (d) $-F_g \sin(\phi)$
- (e) None of the above



Example: Inclined surface



Constant acceleration in 2-D

- Same equations, but treat each direction separately
- x-direction
 - $x_F = x_0 + v_{0x}t_F + \frac{1}{2}a_xt_F^2$

•
$$v_{Fx} = v_{0x} + a_x t_F$$

•
$$v_{Fx}^2 = v_{0x}^2 + 2a_x(\Delta x)$$

- y-direction
 - $y_F = y_0 + v_{0y}t_F + \frac{1}{2}a_yt_F^2$
 - $v_{Fy} = v_{0y} + a_y t_F$
 - $v_{Fy}^2 = v_{0y}^2 + 2a_y(\Delta y)$

 v_{ox} is initial velocity in x

- v_{oy} is initial velocity in y
- a_x is x acceleration
- a_y is y acceleration

Clicker Question 8:

Suppose I throw a ball straight down a 80m cliff with a velocity of 24m/s. What speed does the ball hit the ground with?

- (a) 24.6 m/s
- (b) 28.4 m/s
- (c) 34.5 m/s
- (d) 46.3 m/s
- (e) 57.5 m/s

Clicker Question 9:

Projectiles 1 and 2 are launched at the same time, over level ground with different speeds. Both reach the same height. Which hits the ground first? Ignore air resistance.

- (A) Projectile 1 hits first.
- (B) Projectile 2 hits first.
- (C) They hit at the same time.

(D) There's not enough information to tell.



You want to shoot a ball to a friend of yours standing on a nearby building. There is a picture below of this set-up for your reference. The ball is launched with a speed of 42 m/s, and leaves from a position of 1.5 m above the ground. However, there is a billboard sign between you and the building. You stand such that the ball just barely gets over the billboard at its highest point in its trajectory. (a) At what angle, above the horizontal do you need to shoot the ball for this to happen?



You want to shoot a ball to a friend of yours standing on a nearby building. There is a picture below of this set-up for your reference. The ball is launched with a speed of 42 m/s, and leaves from a position of 1.5 m above the ground. However, there is a billboard sign between you and the building. You stand such that the ball just barely gets over the billboard at its highest point in its trajectory. (b) What is your horizontal distance, d, from the billboard?



You want to shoot a ball to a friend of yours standing on a nearby building. There is a picture below of this set-up for your reference. The ball is launched with a speed of 42 m/s, and leaves from a position of 1.5 m above the ground. However, there is a billboard sign between you and the building. You stand such that the ball just barely gets over the billboard at its highest point in its trajectory. (c) Does the ball strike the roof of the building (you must show work to justify your answer)?



Clicker Question 10:

Suppose a box having a mass M sits on the floor of an elevator that is moving upward and speeding up. Compare the weight of the box (Mg) to the magnitude of the normal force exerted by the elevator floor on the box (F_N) .

(a) $F_N < Mg$ (b) $F_N = Mg$ (c) $F_N > Mg$



Clicker Question 11:

An airplane is flying from San Diego to Los Angeles. Many forces act on the plane, including gravity, air resistance, the trust of the engine, and the lift of the wings. At some point during its trip the velocity of the plane is measured to be constant (which means its altitude is also constant). At this time, the total force on the plane:

- (A) is pointing upward
- (B) is pointing downward
- (C) is pointing forward
- (D) is pointing backward
- (E) is zero